## Infrared Coagulation (IRC) Hemorrhoid Treatment **Preparation Instructions**

## **About Infrared Coagulation:**

Infrared Coagulation (IRC) is mostly performed in an office setting to treat hemorrhoids. It is well tolerated by most patients, and has limited side effects. A small probe contacts the area above the hemorrhoid, exposing the tissue to a burst of infrared light for about a second. This coagulates the veins above the hemorrhoid causing it to shrink and recede. The patient may feel a sensation of heat very briefly, but it is generally not painful. Anesthetic is usually not required.

Please read all the instructions THOROUGHLY!
☐ Please contact our office at <b>303-604-5000</b> if you have questions or concerns.
<mark>2 days</mark> before your exam:
Purchase 1 Plain Fleet Enema from the pharmacy.
☐ If you have a tendency towards constipation, you should purchase 2 enemas.
The day of your exam:
☐ All medications may be taken as usual.
☐ 1 to 1 ½ hours before leaving home: Give yourself the first enema. Hold as long as possible,
up to 20 minutes, and then expel it. Allow yourself to rest for a few minutes. Repeat with second enema if you have a tendency towards constipation, again holding for no more than 20 minutes.
☐ DO NOT EAT OR DRINK ANYTHING AFTER TAKING THE ENEMAS.
*** If you feel the prep is not working, please call our office at 303-604-5000. ***

## **Procedure Cancellation Notice Requirements:**

- Procedure cancellations must be made three (3) business days prior to your appointment.
- Failure to provide the required advance notice will result in a \$50 Total Cancellation Fee.