

# ERCP Preparation Instructions

**Endoscopic Retrograde Cholangiopancreatography (ERCP)** preparation requires a little organization but is an extremely important part of your procedure. **If you do not follow these directions, your ERCP may be cancelled.** Please note, it is your responsibility to contact your insurance company about coverage and cost for your upcoming ERCP procedure. Sometimes, ERCPs result in a cost to the patient. You will need to tell your insurance company to determine if you will have a cost.

## Key Information:

- Bring a list of all your **MEDICATIONS** (including over-the-counter medications), photo ID, insurance card, co-pay/co-insurance/deductible with you to your procedure.
- **DO NOT** bring unnecessary belongings. We are not responsible for lost or misplaced items.
- Stop all clear liquids two hours before your procedure arrival time.

## Transportation on the day of your procedure:

- A responsible adult **MUST BE PRESENT** (18 years or older) with you at check-in before your procedure and **MUST BE ABLE** to pick you up **IMMEDIATELY** when you are discharged. You are **NOT ALLOWED** to drive, take a taxi or bus, or leave the Endoscopy Center alone. If you do not have a responsible driver with you to take you home, your procedure **WILL BE CANCELLED**.
- Because of the sedation, you are **NOT ALLOWED** to use a bus, taxi, Uber/Lyft, Access-a-Ride, walk home, etc. alone.

## Medications:

- **Blood thinners:** including Coumadin (warfarin), Plavix (clopidogrel), Ticlid (ticlopidine hydrochloride), Brilinta (ticagrelor), Aggrastat (tirofiban), Agrylin (anagrelide), Xarelto (rivaroxaban), Pradaxa (dabigatran), Eliquis (apixaban), Lovenox (low molecular weight heparin), Innohep (heparin) or Effient (Prasugrel), You must follow the instructions provided by our nurses.
- **Diabetes/Weight Loss medication:** See instructions below.
- **All other medications:** Including aspirin, should be taken the day of the exam with a sip of water.

## Weight Loss/Diabetes Medications:

- **GLP-1 Agonists:** including Mounjaro (tirzepatide), Zepbound (tirzepatide), Trulicity (dulaglutide), Victoza, Saxenda (liraglutide), Adlyxin (lixisenatide), Ozempic, Wegovy (semaglutide subcutaneous), Bydureon BCise (exenatide extended-release), Byetta (exenatide), Rybelsus (semaglutide tablet), Stop the oral or injectable medication one week prior to your procedure.
- **Anorectics:** including Adipex-P, Lomaira, Suprenza (phentermine hydrochloride), Qsymia (phentermine hydrochloride/topiramate), Stop 5 days prior to your procedure.

## If you have diabetes, follow these instructions:

- **Long-Acting Insulins** (ex. Lantus, Basaglar, Semglee, Toujeo, Insulin Glargine): Decrease basal rate by  $\frac{1}{2}$  the day before and the day of your procedure.
- **Rapid-Acting and Regular Short-Acting Insulin** (ex. Lispro, Humalog, Admelog, Regular Insulin): If your blood sugar is high and you need to take a correction, take  $\frac{1}{2}$  or 50% of your usual correction. This applies to the day before and the day of your procedure.
- **Insulin Pump:** Decrease basal rate by  $\frac{1}{2}$  the day before and the day of your procedure. Do not administer bolus doses.
- **Other Diabetic Oral & Injectable Medications** (ex. Metformin, Glipizide, Sulfonylureas, Tradjenta, Dipeptidyl Peptidase IV (DPP IV) Inhibitors, Januvia, Janumet, Sodium-Glucose Cotransporter-2 (SGLT2) Inhibitors): Do not take the day before or the day of your procedure.
- **Check your blood sugar at same intervals as usual, and additionally, if needed:** If your blood sugar is less than 60, please take glucose tablets at any time before your procedure with a sip of water.

# Checklist to Ensure Proper Preparation

To ensure the proper completion of your ERCP preparation, we strongly suggest that you **PRINT OUT** this document and check off each of the instructions as you complete them. This will greatly help you in following the detailed instructions.

## 5 days before your procedure:

- DO NOT** take products that contain iron such as multi-vitamins or iron supplements.

## Day of your procedure:

- DO NOT eat solid food for 9 hours before your arrival time.**  
**For example, if your arrival time is at 9 AM, do NOT eat after MIDNIGHT.**
- Discontinue** mints, gum, hard candy, and chewing tobacco 9 hours before your arrival time.
- Follow a **clear liquid diet** using the charts below:

### CLEAR LIQUIDS - YOU CAN DRINK:

Gatorade, Pedialyte, Powerade, Kool-Aid
Gatorade G2, Propel, Crystal Light
Popsicles
Jell-o, Juice without Pulp (ex. Apple Juice)
Black Coffee
Tea, Water
Clear Sodas (ex. Sprite, Ginger Ale)
Sparkling water

### DO NOT DRINK:

<b>NO</b> Red or Purple Liquids
<b>NO</b> Milk or Non-dairy Creamers
<b>NO</b> Alcohol
<b>NO</b> Broth or Soup
<b>NO</b> Juice with Pulp (ex. pineapple juice)
<b>NO</b> Liquid you cannot see through
<b>NO</b> Apple Sauce
<b>NO</b> Smoothies

- Morning medications may be taken as usual except for above-mentioned medications.**
- No marijuana on the day of your procedure.**
- If you have **diabetes**, please see Diabetes instructions on page 1.

## 2 hours - before your procedure check-in time

- Discontinue** mints, gum, hard candy, and chewing tobacco.
- 2 hours before check-in: STOP DRINKING ALL LIQUIDS!**

**Congratulations, you have completed your prep!**

### Procedure Cancellation Notice Requirements:

- Procedure cancellations must be made **three (3) business days prior to your appointment.**
- Failure to provide the required advance notice will result in a **\$150 Total Cancellation Fee.** (\$50 Physician cancellation fee from Gastroenterology of the Rockies + \$100 Facility cancellation fee from the endoscopy center).