

# Suprep Bowel Preparation Instructions

**These instructions are for patients WITHOUT severe kidney and heart disease. If you have severe kidney or heart disease, please call our office for alternative instructions: 303-604-5000**

This bowel preparation requires a little organization but is an extremely important part of your colonoscopy to ensure you have the highest quality exam. Bowel preparation starts days prior to your procedure, as you need to stop taking certain medications or supplements and change your diet. **If you do not follow these directions, your colonoscopy may be cancelled.**

## Key Information:

- **DO NOT EAT** any solid food after 7pm two days before your colonoscopy. Drink only clear fluids.
- Bring a list of all your **MEDICATIONS** (including over-the-counter medications), photo ID, insurance card, co-pay/co-insurance/deductible with you to your procedure.
- **DO NOT** bring unnecessary belongings. We are not responsible for lost or misplaced items.
- Stop all clear liquids three hours before your procedure arrival time.
- Your colon must be **COMPLETELY EMPTY** so that your doctor can clearly view your colon. Follow all instructions in this handout **EXACTLY** as they are written. If you have completed your prep, and your stools are not clear or yellow, please contact our office at 303-604-5000, even if after hours.

## Transportation on the day of your procedure:

- A responsible adult **MUST BE PRESENT** (18 years or older) with you at check-in before your procedure and **MUST BE ABLE** to pick you up **IMMEDIATELY** when you are discharged. You are **NOT ALLOWED** to drive, take a taxi or bus, or leave the Endoscopy Center alone. If you do not have a responsible driver with you to take you home, your procedure **WILL BE CANCELLED**.
- Because of the sedation, you are **NOT ALLOWED** to use a bus, taxi, Uber/Lyft, Access-a-Ride, walk home, etc. alone.

## Medications:

- **Blood thinners:** including Coumadin (warfarin), Plavix (clopidogrel), Ticlid (ticlopidine hydrochloride), Brilinta (ticagrelor), Aggrastat (tirofiban), Agrylin (anagrelide), Xarelto (rivaroxaban), Pradaxa (dabigatran), Eliquis (apixaban), Lovenox (low molecular weight heparin), Innohep (heparin) or Effient (Prasugrel), You must follow the instructions provided by our nurses.
- **Diabetes/Weight Loss medication:** See instructions below.
- **ACE Inhibitor Blood Pressure medications:** Do NOT take medications ending 'pril' the day of your procedure. Examples: lisinopril, benazepril, ramipril, captopril, enalapril, quinapril
- **ARB Blood Pressure medications:** Do NOT take medications ending in 'sartan' the day of your procedure. Examples: losartan, irbesartan, candesartan, valsartan, telmisartan, olmesartan
- **Diuretics:** Do NOT take "water" pills the day of your procedure. Examples: furosemide (Lasix), bumetanide (Bumex), torsemide, hydrochlorothiazide (HCTZ), spironolactone (Aldactone), indapamide, chlorothiazide, chlorthalidone, metolazone
- **All other medications:** Including aspirin, should be taken the day of the exam with a sip of water.

## Weight Loss/Diabetes Medications:

- **GLP-1 Agonists (Injectable):** including Mounjaro (tirzepatide), Zepbound (tirzepatide), Trulicity (dulaglutide), Victoza, Saxenda (liraglutide), Adlyxin (lixisenatide), Ozempic (semaglutide), Wegovy (semaglutide), Bydureon BCise (exenatide), Byetta (exenatide), Stop one week prior to your procedure.
- **GLP-1 Agonists (Oral):** including Rybelsus (semaglutide tablet), Stop 24 hours prior to procedure.
- **Anorectics:** including Adipex-P, Lomaira, Suprenza (phentermine hydrochloride), Qsymia (phentermine hydrochloride/topiramate), Stop 5 days prior to your procedure.

**If you have diabetes, follow these instructions:**

- **Long-Acting Insulins** (ex. Lantus, Basaglar, Semglee, Toujeo, Insulin Glargine): Decrease basal rate by ½ the day before and the day of your procedure.
- **Rapid-Acting and Regular Short-Acting Insulin** (ex. Lispro, Humalog, Admelog, Regular Insulin): If your blood sugar is high and you need to take a correction, take ½ or 50% of your usual correction. This applies to the day before and the day of your procedure.
- **Insulin Pump**: Decrease basal rate by ½ the day before and the day of your procedure. Do not administer bolus doses.
- **SGLT2 Inhibitors**: including Jardiance (empagliflozin), Brenzavvy (bexagliflozin), Glyxambi (empagliflozin/linagliptin), Invokamet (canagliflozin/metformin), Invokana (canagliflozin), Qtern (dapagliflozin/saxagliptin), Segluromet (ertugliflozin/metformin), Steglatro (ertugliflozin), Steglujan (ertugliflozin/sitagliptin), Synjardy (empagliflozin/metformin), Inpefa (sotagliflozin), Trijardy (empagliflozin/linagliptin/metformin), Xigduo (dapagliflozin/metformin), Farxiga (dapagliflozin), Stop these medications four days prior to your procedure.
- **Other Diabetic Oral & Injectable Medications**: including Glucophage (metformin), Glucotrol/Glynase Pres Tab/Micronase (glipizide), Amaryl (glimepiride), DiaBeta (glyburide), Tradjenta (linagliptin), Januvia (sitagliptin), Nesina (alogliptin), Onglyza (saxagliptin), Zituvio (sitagliptin), Actos (pioglitazone), Avandia (rosiglitazone) or any combination of these medications, Stop these medications the day before your procedure.
- **Check your blood sugar at same intervals as usual, and additionally, if needed**: If your blood sugar is less than 60, please take glucose tablets at any time before your procedure with a sip of water.

## Checklist to Ensure Proper Bowel Preparation

To ensure the proper completion of your colonoscopy preparation, we strongly suggest that you **PRINT OUT** this document and check off each of the instructions as you complete them. This will greatly help you in following the detailed instructions and help ensure you receive a proper colonoscopy.

**GROCERY STORE LIST**: Purchase the items below at least five days before your procedure:

- ☐ **1** - Suprep Bowel Kit
- ☐ **4** - Gas-X tablets (simethicone).
- ☐ **1** - 16-gram packet of magnesium citrate OR 10 oz bottle of magnesium citrate

**5 days** before your colonoscopy:

- ☐ **Make sure you have purchased your bowel preparation items!** (listed above!)
- ☐ **DO NOT** take medications that stop diarrhea such as Imodium (loperamide), Pepto-Bismol (bismuth subsalicylate) or Lomotil (diphenoxylate/atropine).
- ☐ **DO NOT** take fiber supplements such as Metamucil, Citrucel, Psyllium or Benefiber.
- ☐ **DO NOT** take products that contain iron such as multi-vitamins or iron supplements.

**3 days** before your colonoscopy:

- ☐ **Discontinue high-fiber foods** – Please see Low Fiber Diet tips attached.

## Checklist: Two Days Before Your Colonoscopy

- ☐ **DO NOT** eat solid food after 7 pm until after your procedure.
- ☐ Follow a **clear liquid diet** using the charts below:

### CLEAR LIQUIDS - YOU CAN DRINK:

Gatorade, Pedialyte, Powerade, Kool-Aid
Gatorade G2, Propel, Crystal Light
Popsicles
Jell-o, Juice without Pulp (ex. Apple Juice)
Black Coffee
Tea, Water
Clear Sodas (ex. Sprite, Ginger Ale)
Sparkling water

### DO NOT DRINK:

<b>NO</b> Red or Purple Liquids
<b>NO</b> Milk or Non-dairy Creamers
<b>NO</b> Alcohol
<b>NO</b> Broth or Soup
<b>NO</b> Juice with Pulp (ex. pineapple juice)
<b>NO</b> Liquid you cannot see through
<b>NO</b> Apple Sauce
<b>NO</b> Smoothies

## Checklist: The Day Before Your Colonoscopy

- ☐ If you have **diabetes**, please see diabetes instructions above.
- ☐ Only drink clear liquids the **ENTIRE DAY** before your procedure. **NO solid foods**.
- ☐ Drink at least 8-ounces of **clear liquids EVERY HOUR** after waking up. See charts above.

### 5:00 pm - the evening before your procedure:

- ☐ Take two tablets of simethicone/Gas X

### 6:00 pm - the evening before your procedure:

- ☐ Pour one 6-ounce bottle of Suprep liquid into the mixing container.
- ☐ Add cool water to the 16-ounce line on the container and mix.
- ☐ Drink all the liquid in the container.
- ☐ Drink two more 16-ounce glasses of water over the next hour (one hour).
- ☐ **Before going to sleep:** Take two additional simethicone tablets.
- ☐ **Set your alarm!** Make sure to wake up at least **5 hours** before your procedure check-in time.

### 9:00 pm - the evening before your procedure:

- ☐ If you have not had any bowel movements, drink either the 10 oz bottle of magnesium citrate or a 16-gram packet of magnesium citrate (mixed with 8 oz of clear liquid of your choice).

# Checklist: The Day of Your Colonoscopy

- ☐ If you have **diabetes**, please see diabetes instructions above.
- ☐ **No marijuana.**
- ☐ **No mints, gum, hard candy or chewing tobacco.**
- ☐ You may drink clear liquids up to 3 hours before your procedure check-in time. **NO solid foods.**

**5 hours - before your procedure check-in time** - this may be in the middle of the night or early AM:

- ☐ Pour one 6-ounce bottle of Suprep liquid into the mixing container.
- ☐ Add cool water to the 16-ounce line on the container and mix.
- ☐ Drink all the liquid in the container.
- ☐ Drink two more 16-ounce glasses of water over the next hour (one hour).

**3 hours - before your procedure check-in time**

- ☐ **3 hours before check-in: STOP DRINKING ALL LIQUIDS!**

Please note, all patients of childbearing age with a uterus will be asked to submit a urine sample prior to the procedure to test for pregnancy.

**Congratulations, you have completed your prep!** Please bring your medication list (including over-the-counter medications), photo ID, insurance card, co-pay/co-insurance/deductible with you to your procedure.

If you have any questions throughout your prep, please see our procedure FAQ's:  
<https://www.gastrorockies.com/procedure-faqs/>

## Procedure Cancellation Notice Requirements:

- Procedure cancellations must be made **three (3) business days prior to your appointment.**
- Failure to provide the required advance notice will result in a **\$150 Total Cancellation Fee.** (\$50 Physician cancellation fee from Gastroenterology of the Rockies + \$100 Facility cancellation fee from the endoscopy center).

# Low Fiber Diet

A low fiber diet limits the amount of food waste that must move through the large intestine.

## Approved Foods

### Bread, cereal, rice, and pasta

- White bread, rolls, biscuits, croissants, melba toast
- Waffles, French toast, pancakes
- White rice, noodles, pasta, and macaroni
- Peeled cooked potatoes
- Plain crackers, saltines
- Farina, cream of rice
- Puffed rice, rice krispies, corn flakes, special K

### Meat

- Ground (beef, chicken, turkey)
- Lamb
- Pork
- Veal
- Fish
- Poultry

### Vegetables

- Cooked carrots
- Asparagus tips
- Green or wax beans
- Pumpkin
- Lima Beans

### Fruits

- Bananas, Honeydew, cantaloupe

### Milk/Dairy

- Milk-plain or flavored, eggs
- Yogurt, custard, ice cream, cheese, cottage cheese

### Fats, snacks, sweets, condiments, and beverages

- Margarine, butter, oils
- Mayonnaise
- Sour cream
- Dressing
- Plain gravy
- Sugar
- Clear jelly
- Honey
- Syrup
- Spices/cooked herbs
- Bouillon, broth
- Coffee, tea, carbonated drinks
- Plain cake, cookies
- Gelatin, plain pudding, popsicles
- Hard candy or pretzels
- Mustard, ketchup

## Foods to Avoid

- Breads or rolls with nuts, seeds or fruit: whole wheat, pumpernickel, rye, corn bread, chia seeds
- Brown or wild rice, buckwheat, quinoa
- Vegetables (raw or steamed, with seeds) - winter squash, peas, broccoli, Brussels sprouts
- Cabbage, onions, cauliflower, sprouts, baked beans, peas, corn, potatoes with skin
- Sauerkraut
- Fruits (raw or dried) - all berries, figs, dates/raisins, prunes, prune juice
- Yogurt with nuts or seeds
- Processed meat - hotdogs, sausage, cold cuts, tough meat with gristle