

# Clenpiq Bowel Preparation Instructions

**These instructions are for people who do not have serious kidney or heart problems. If you have kidney or heart disease, please call our office: 303-604-5000.**

Cleaning out your colon helps your doctor see clearly during your colonoscopy. If your colon is not clean, your test may be **canceled**, and you may have to **reschedule**.

## Key Information:

- **Read these instructions at least two weeks before your procedure.**
- **Stop eating solid food at 7 PM, two days before your colonoscopy.**
- Drink **only clear liquids** after that.
- Stop drinking all liquids **3 hours before your check-in time.**
- Diarrhea is expected during your prep.
- Nausea is common. If you are prone to nausea, please call the office **during regular business hours** to speak with a nurse about whether a prescription to help is needed. Please see FAQs for common questions about what to expect and tips. Scan this QR Code or paste this link in your browser:  
<https://www.gastrorockies.com/procedure-faqs/>
- **Print these instructions and check off each step as you do it.**  
It is important to follow all these instructions to ensure that the physician can safely visualize your colon during the procedure.
- If you can become pregnant and have a uterus, you will need to give a urine sample before your procedure to check for pregnancy or sign a waiver. Pregnant patients will be canceled.
- **Do you have constipation and/or infrequent bowel movements?** You should not be backed up going into the prep. Contact office if this is a concern for you to adjust your prep.



## Transportation on the day of your procedure:

- You must bring an adult (18+) with you. They need to stay in the facility the whole time and take you home after your procedure.
- **You cannot go home alone** (no walking, bus, taxi, Uber, etc.). If you don't have someone to take you, your procedure will be **canceled**.
- You will be at the facility for approximately 2 hours, and your driver is required to check in with you and stay at the facility until you are discharged.

## Medication Information:

If you take medications for blood pressure, weight loss, diabetes, blood thinning or kidney issues, please review this section for medications that must be held prior to your procedure. If these medications are not held according to these instructions, your procedure will be canceled.

- **Blood thinners:** including Coumadin (warfarin), Plavix (clopidogrel), Ticlid (ticlopidine hydrochloride), Brilinta (ticagrelor), Aggrastat (tirofiban), Agrylin (anagrelide), Xarelto (rivaroxaban), Pradaxa

(dabigatran), Eliquis (apixaban), Lovenox (low molecular weight heparin), Innohep (heparin) or Effient (Prasugrel), you must follow the instructions provided by our nurses.

- **ACE Inhibitor Blood Pressure medications:** Do NOT take medications ending 'pril' the day of your procedure. Examples: lisinopril, benazepril, ramipril, captopril, enalapril, quinapril
- **ARB Blood Pressure medications:** Do NOT take medications ending in 'sartan' the day of your procedure. Examples: losartan, irbesartan, candesartan, valsartan, telmisartan, olmesartan
- **Diuretics:** Do NOT take "water" pills the day of your procedure. Examples: furosemide (Lasix), bumetanide (Bumex), torsemide, hydrochlorothiazide (HCTZ), spironolactone (Aldactone), indapamide, chlorothiazide, chlorthalidone, metolazone

#### GLP-1 Agonists and Other Weight Loss Medications:

- **GLP-1 Agonists (Injectable):** including Mounjaro (tirzepatide), Zepbound (tirzepatide), Trulicity (dulaglutide), Victoza, Saxenda (liraglutide), Ozempic (semaglutide), Wegovy (semaglutide), Bydureon BCise (exenatide), Byetta (exenatide), stop one week prior to your procedure.
- **GLP-1 Agonists (Oral):** including Rybelsus (semaglutide tablet), stop 24 hours prior to procedure.
- **Anorectics:** including Adipex-P, Lomaira, Suprenza (phentermine hydrochloride), Qsymia (phentermine hydrochloride/topiramate), stop 5 days prior to your procedure.

#### Insulin and Blood Sugar Checking Instructions:

- **Long-Acting Insulins** (ex. Lantus, Basaglar, Semglee, Toujeo, Soliqua, Insulin Glargine): Decrease basal rate by  $\frac{1}{2}$  the day before and the day of your procedure.
- **Rapid-Acting and Regular Short-Acting Insulin** (ex. Lispro, Humalog, Admelog, Regular Insulin): If your blood sugar is high and you need to take a correction, take  $\frac{1}{2}$  or 50% of your usual correction. This applies to the day before and the day of your procedure.
- **Insulin Pump:** Decrease basal rate by  $\frac{1}{2}$  the day before and the day of your procedure. Do not administer bolus doses.
- **Check your blood sugar at same intervals as usual, and additionally, if needed:** If your blood sugar is less than 60, please take glucose tablets at any time before your procedure with a sip of water.

#### Other Medications that Lower Blood Sugar Instructions:

- **SGLT2 Inhibitors:** including Jardiance (empagliflozin), Brenzavvy (bexagliflozin), Glyxambi (empagliflozin/linagliptin), Invokamet (canagliflozin/metformin), Invokana (canagliflozin), Qtern (dapagliflozin/saxagliptin), Segluromet (ertugliflozin/metformin), Steglatro (ertugliflozin), Steglujan (ertugliflozin/sitagliptin), Synjardy (empagliflozin/metformin), Inpefa (sotagliflozin), Trijardy (empagliflozin/linagliptin/metformin), Xigduo (dapagliflozin/metformin), Farxiga (dapagliflozin), stop these medications 4 days prior to your procedure.
- **Other Medicines that Help Control Blood Sugar:** including Glucophage (metformin), Glucotrol/Glynase Pres Tab/Micronase (glipizide), Amaryl (glimepiride), DiaBeta (glyburide), Tradjenta (linagliptin), Januvia (sitagliptin), Nesina (alogliptin), Onglyza (saxagliptin), Zituvio (sitagliptin), Actos (pioglitazone), Avandia (rosiglitazone) or any combination of these medications, stop these medications the day before your procedure.

#### Other Medication Instructions:

- **Stop medications** for diarrhea such as Imodium (loperamide), Pepto-Bismol (bismuth subsalicylate) or Lomotil (diphenoxylate/atropine) 5 days before your procedure
- **Stop iron products** like vitamins with iron or iron pills 5 days before your procedure
- **Stop fiber supplements** like Metamucil, Citrucel, Psyllium or Benefiber 5 days before your procedure
- **All other medications:** Including aspirin, should be taken the day of the procedure with a sip of water

# Checklist to Ensure Proper Bowel Preparation

**Prep Items Needed:** Purchase the items below at least five days before your procedure:

- ☐ **1** - Clenpiq Bowel Kit
- ☐ **4** - Gas-X tablets containing 80 mg or 125 mg of simethicone each.
- ☐ **1** - Easily measurable 8-ounce glass/cup: You will be drinking multiple quantities of 8 oz liquids.
- ☐ **1** - 16-gram packet of magnesium citrate OR 10 oz bottle of magnesium citrate

**3 days** before your colonoscopy:

- ☐ **Discontinue high-fiber foods** – Please see Low Fiber Diet tips attached.

## Checklist: Two Days Before Your Colonoscopy

- ☐ **DO NOT eat solid food after 7 pm until after your procedure.**  
**Example:** If your procedure is on Monday, stop eating solid food at 7pm on Saturday evening.
- ☐ Follow a **clear liquid diet** using the charts below:

### CLEAR LIQUIDS - YOU CAN DRINK:

|                                              |
|----------------------------------------------|
| Gatorade, Pedialyte, Powerade, Kool-Aid      |
| Gatorade G2, Propel, Crystal Light           |
| Popsicles                                    |
| Jell-o, Juice without Pulp (ex. Apple Juice) |
| Black Coffee                                 |
| Tea, Water                                   |
| Clear Sodas (ex. Sprite, Ginger Ale)         |
| Sparkling water                              |

### DO NOT DRINK:

|                                                 |
|-------------------------------------------------|
| <b>NO</b> Red or Purple Liquids                 |
| <b>NO</b> Milk or Non-dairy Creamers            |
| <b>NO</b> Alcohol                               |
| <b>NO</b> Broth or Soup                         |
| <b>NO</b> Juice with Pulp (ex. pineapple juice) |
| <b>NO</b> Liquid you cannot see through         |
| <b>NO</b> Apple Sauce                           |
| <b>NO</b> Smoothies                             |

# Checklist: The Day Before Your Colonoscopy

- ☐ Only drink clear liquids the **ENTIRE DAY** before your procedure. **NO solid foods.**
- ☐ Drink at least 8-ounces of **clear liquids EVERY HOUR** after waking up.

## 5:00 pm - the evening before your procedure:

- ☐ Take two tablets of simethicone/Gas X

## 6:00 pm - the evening before your procedure:

- ☐ Drink one bottle of Clenpiq.
- ☐ Over the next five hours, drink at least 5 cups (8 ounces each) of clear liquid at your own pace.
- ☐ **Before going to sleep:** Take two additional simethicone tablets.
- ☐ **Set your alarm!** Make sure to wake up at least **5 hours** before your procedure check-in time.

## 9:00 pm - the evening before your procedure:

- ☐ If you haven't had a bowel movement yet, drink a 10 oz bottle of magnesium citrate or mix a 16-gram packet of magnesium citrate with 8 oz of clear liquid and drink it.

# Checklist: The Day of Your Colonoscopy

- ☐ No marijuana.
- ☐ No mints, gum, hard candy, nicotine pouches/patches or chewing tobacco.
- ☐ You may drink clear liquids up to 3 hours before your procedure check-in time. **NO solid foods.**

**5 hours - before your procedure check-in time** - this may be in the middle of the night or early AM:

- ☐ Drink one bottle of Clenpiq.
- ☐ Then drink one cup (8 ounces each) of clear liquid, every 15 minutes for at least four cups. **If your stools are not light yellow/clear and you did not already drink the magnesium citrate after the first half of the prep, drink a 10 oz bottle of magnesium citrate or 16-gram packet of magnesium citrate with 8 oz of clear liquid.**

**3 hours - before your procedure check-in time**

- ☐ **3 hours before check-in: STOP DRINKING ALL LIQUIDS!**

**\*\*\*By this point, your stool should look light yellow and clear—similar to urine.** If it is still brown, cloudy or has any solid pieces, your colon may not be clean enough for your procedure. Please call our office at 303-604-5000 **at 6:30 am for further instructions\*\*\***

**You're all done with your prep—great job!**

**Bring these things with you to your procedure:**

- ☐ A list of all your medicines (even ones you buy without a prescription)
- ☐ A photo ID
- ☐ Your insurance card
- ☐ Payment card for your co-pay, co-insurance or deductible
- ☐ Don't bring valuables—we are not responsible for lost items.
- ☐ A driver who can stay at the facility during your entire visit and drive you home

## **Procedure Cancellation Notice Requirements:**

- Procedure cancellations must be made **seven (7) business days prior to your appointment.**
- Failure to provide the required advance notice will result in a **\$300 Cancellation Fee.**

## ✓ Low-Fiber Foods to Eat (Safe 3 Days Before Colonoscopy)

| Category                        | Recommended Foods                                                                                                                                           |
|---------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Milk &amp; Milk Products</b> | Buttermilk, Custard, Evaporated/Powdered Milk, Low-fat Cheese, Low-fat Ice Cream, Skim/Low-fat Milk, Sherbet, Smooth Yogurt (no seeds/berries/nuts)         |
| <b>Vegetables</b>               | Well-cooked veggies without seeds (e.g., carrots, spinach), Lettuce, Potatoes (no skin), Strained Veggie Juice                                              |
| <b>Meats &amp; Protein</b>      | Eggs, Smooth Nut Butters (e.g., creamy peanut butter), Tofu, Tender/Well-cooked Meat, Poultry, Fish                                                         |
| <b>Grains</b>                   | White Bread, White Rice, Pasta, Crackers, Cereals made with refined flour, Cooked cereals (farina, creamy rice)                                             |
| <b>Fruits</b>                   | Juice without pulp (except prune), Peeled Apple, Ripe Banana, Melons, Soft/Pureed canned fruit (except pineapple)                                           |
| <b>Snacks &amp; Sweets</b>      | Jell-O, Hard Candy, Plain Cake/Cookies, Plain Pudding, Low-fat Ice Cream, Pretzels, Sherbet, Popsicles                                                      |
| <b>Condiments</b>               | Ketchup, Mustard, Butter, Margarine, Oils, Mayo, Sour Cream, Plain Gravy, Clear Jelly, Honey, Syrup, Broth, Bouillon, Cooked Herbs, Allowed Vegetable Soups |

## ✗ High-Fiber Foods to Avoid (3 Days Before Colonoscopy)

| Category                        | Avoid These Foods                                                                                                                                                                                       |
|---------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Milk &amp; Milk Products</b> | Yogurt with seeds, berries, or nuts                                                                                                                                                                     |
| <b>Vegetables</b>               | Broccoli, Cabbage, Cauliflower, Corn, Fried Vegetables, Greens (mustard, collards, etc.), Mushrooms, Onions, Okra, Peppers, Raw Veggies (except lettuce), Potato Skins, Brussels Sprouts, Winter Squash |
| <b>Meats &amp; Protein</b>      | Chunky Nut Butters, Dried/Legume Beans (peas, lentils), Nuts/Seeds, Tough or Chewy Meats, Processed meat (hot dogs, sausage, cold cuts)                                                                 |
| <b>Grains</b>                   | Brown/Wild Rice, Whole-grain Cereals, Whole-Wheat or Seeded Breads/Rolls/Pasta/Crackers                                                                                                                 |
| <b>Fruits</b>                   | Raw Fruits (except peeled apple, ripe banana, melon), Dried Fruits (e.g., raisins), Prunes & Prune Juice, Canned Berries/Cherries                                                                       |
| <b>Snacks/Sweets/Condiments</b> | Popcorn, Candy with Nuts/Seeds, Desserts with Whole Grains, Dried Fruit, Nuts or Coconut, Jam, Marmalade, Pickles, Olives, Relish, Horseradish, Sauerkraut                                              |